Food Allergy Action Plan

Allergy to:			Camp Swamp
Weight: lbs. Asthma: ☐ Yes (higher risk for a severe reaction) ☐ No			
Extremely reactive to the following foods: THEREFORE: ☐ If checked, give epinephrine immediately for ANY symptoms if the allergen was likely eaten. ☐ If checked, give epinephrine immediately if the allergen was definitely eaten, even if no symptoms are noted.			
ingestion: One or more LUNG: HEART: THROAT: MOUTH: SKIN:	Pale, blue, faint, weak pulse, dizzy, confused Tight, hoarse, trouble breathing/swallowing	asthma	ering (see box all medications:* ne inchodilator) if halers/bronchodilators ded upon to treat a
MILD SYMPT MOUTH: SKIN: GUT:	Itchy mouth A few hives around mouth/face, mild itch Mild nausea/discomfort	parent 3. If symptoms above), USE 4. Begin monitor	ident; alert rofessionals and
Medications/Doses below)			
Epinephrine (brand and dose):Antihistamine (brand and dose):			
Other (e.g., inhaler-bronchodilator if asthmatic):			
Monitoring Stay with student; alert healthcare professionals and parent. Tell rescue squad epinephrine was given; request an ambulance with epinephrine. Note time when epinephrine was administered. A second dose of epinephrine can be given 5 minutes or more after the first if symptoms persist or recur. For a severe reaction, consider keeping student lying on back with legs raised. Treat student even if parents cannot be reached. See back/attached for auto-injection technique.			
Parent/Guardian Signature Date Physician/Healthcare Provider Signature Date			